arXiv:math/0702137v2 [math.RT] 13 Feb 2007

Bilinear forms on $\mathfrak{sl}_2$-modules and a hypergeometric identity

Johan Kåhrström

February 2, 2008

Abstract

In this paper we study properties of a certain bilinear form on finite dimensional $\mathfrak{sl}_2(\mathbb{R})$-modules, and how these properties behave with respect to tensor products of modules. An attempt to determine the signature of this form leads to an interesting identity for the hypergeometric series ${}_3F_2$, which is known as the Karlsson-Minton identity.

1 Introduction

Let X, Y, H be the standard basis of the Lie-algebra $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{R})$. These elements satisfy the following defining relations:

$$[X,Y] = H, \qquad [H,X] = 2X, \qquad [H,Y] = -2Y. \tag{1}$$

Furthermore, let $*$ be the anti-involution on $\mathfrak{g}$ given by

$$X^* = X, \qquad Y^* = Y, \qquad H^* = -H. \tag{2}$$

Let V be a finite dimensional vector space over $\mathbb{R}$, and Q be a non-degenerate symmetric bilinear form on V. A *$*$-representation* of $\mathfrak{g}$ is a representation $\varphi : \mathfrak{sl}_2(\mathbb{R}) \to \mathfrak{gl}(V)$ such that

$$Q\big(\varphi(x)u, v\big) = Q\big(u, \varphi(x^*)v\big)$$

for all $x \in \mathfrak{g}$, $u, v \in V$. We will say that V is a $*$-representation of $\mathfrak{g}$ with respect to Q, and write xv instead of $\varphi(x)v$. Given two $*$-representations U and V, with respect to the bilinear forms Q and R respectively, the tensor product $U \otimes V$ is also a $*$-representation, with respect to the induced bilinear form $Q \otimes R$.

For $m \in \{0, 1, 2, \dots\}$, denote by V_m the $(m+1)$-dimensional irreducible representation of $\mathfrak{g}$, which has the basis $\{e_{-m+2i} \mid 0 \leq i \leq m\}$, and where the action of our generators in this basis is given by the following formulae (for simplicity, we assume $e_{-m-2} = e_{m+2} = 0$), see [FH, Section 11],

$$\begin{aligned} He_{m-2i} &= ie_{m-2i}, \\ Ye_{m-2i} &= e_{m-2(i+1)}, \\ Xe_{m-2i} &= i(m-i+1)e_{m-2(i-1)}. \end{aligned} \tag{3}$$

Let $m, n \in \{0, 1, 2, \dots\}$. Let further Q and R be non-degenerate symmetric bilinear forms on V_m and V_n respectively, which make these representations of $\mathfrak{g}$ into $*$-representations.

For $0 \leq k \leq \min\{m, n\}$, let $P_{-m-n+2k}$ denote the kernel of Y when acting on the weight subspace of $V_m \otimes V_n$ of weight $-m-n+2k$ (which is by definition the set of all $v \in V_m \otimes V_n$ such that $Hv = (-m-n+2k)v$). The form $Q \otimes R$ induces the following symmetric bilinear form on $P_{-m-n+2k}$:

$$\begin{array}{rcl} \omega_k : P_{-m-n+2k} \times P_{-m-n+2k} & \to & \mathbb{R} \\ (v, w) & \mapsto & Q \otimes R(v, X^{m+n-2k} w) \end{array} \tag{4}$$

The following question was formulated by K.-H. Fieseler in connection to the study of Hodge-Riemann relations for polytopes, see [BBFK].

Question: *Are the forms ω_k positive/negative definite, alternating with k, $0 \leq k \leq \min\{m, n\}$?*

In [BBFK], this question is answered using some abstract arguments. In this paper, we propose a brute-force solution based on explicit calculation. The advantage of the latter method is that it leads to an identity for the hypergeometric series ${}_3F_2$, which is known as the Karlsson-Minton identity [M, K]. We refer the reader to [PWZ] for a survey on hypergeometric identities. It is interesting to see such an identity arise from representation theory of Lie algebras.

This is the second draft of this paper, and when writing the first draft I was not aware of the previous occurence of the identity in [M, K]. I would like to thank Hjalmar Rosengren and Michael Schlosser for pointing this out to me.

In Section 2, we show how the positive answer to the above question reduces to a ${}_3F_2$ identity. This essentially amounts to determining the sign of certain Clebsch-Gordan coefficients of the tensor product $V_m \otimes V_n$. For a more general treatment of determining Clebsch-Gordan coefficiets, see [VK, Chapter 8].

2 Solution of the problem

Let $m \in \{0, 1, 2, \dots\}$ and $Q : V_m \times V_m \to \mathbb{R}$ be a non-degenerate symmetric bilinear form such that V_m becomes a $*$-representation with respect to Q.

Lemma 1. *The bilinear form Q satisfies the following:*

- *(i)* $Q(e_i, e_j) = 0$ *for all* $i \neq -j \in \{-m, -m+2, \dots, m-2, m\}$.
- *(ii)* $Q(e_i, e_{-i}) = Q(e_j, e_{-j})$ *for all* $i, j \in \{-m, -m+2, \dots, m-2, m\}$.

Proof. From (2) we get that

$$iQ(e_i, e_j) = Q(He_i, e_j) = Q(e_i, -He_j) = -jQ(e_i, e_j)$$

for all $i, j \in \{-m, -m+2, \dots, m-2, m\}$, so $Q(e_i, e_j) = 0$ provided that $i \neq -j$. This proves (i).

Furthermore,

$$Q(e_i, e_{-i}) = Q(Ye_{i+2}, e_{-i}) = Q(e_{i+2}, Ye_{-i}) = Q(e_{i+2}, e_{-i-2})$$

for all $i \in \{-m, -m+2, \dots, m-2\}$. This proves (ii).

□

Recall from [FH, Section 11] that for $m, n \in \{0, 1, 2, \dots\}$ we have

$$V_m \otimes V_n = V_{|m-n|} \oplus V_{|m-n|+2} \oplus \cdots \oplus V_{m+n}. \tag{5}$$

In particular, we can depict the explicit basis and subspaces P_i's in the module $V_m \otimes V_n$ as follows:



To distinguish the bases of V_m and V_n we assume that the set $\{e_{-m}, e_{-m+2}, \dots, e_m\}$ is a basis of V_m and $\{\tilde{e}_{-n}, \tilde{e}_{-n+2}, \dots, \tilde{e}_n\}$ is a basis of V_n. Let Q and R be symmetric bilinear forms on V_m and V_n respectively, which make these representations into $*$-representations. By Lemma 1, there exist $q, r \in \mathbb{R}$ such that $Q(e_i, e_{-i}) = q$ and $R(\tilde{e}_i, \tilde{e}_{-i}) = r$ for all appropriate i.

The subspace of $V_m \otimes V_n$ of weight $-m-n+2k$ is spanned by the vectors

$$e_{-m} \otimes \tilde{e}_{-n+2k},\ e_{-m+2} \otimes \tilde{e}_{-n+2(k-1)},\ \ldots,\ e_{-m+2k} \otimes \tilde{e}_{-n}. \tag{6}$$

Lemma 2. *For each* $0 \leq k \leq \min\{m, n\}$ *the vector*

$$b_{-m-n+2k} = \sum_{i=0}^{k} (-1)^i e_{-m+2i} \otimes \tilde{e}_{-n+2(k-i)}.$$

forms a basis of the vector space $P_{-m-n+2k}$.

Proof. From the decomposition (5) it follows that $\dim P_{-m-n+2k} = 1$. Hence, to find a basis of $P_{-m-n+2k}$ it is sufficient to find a non-zero element of weight $-m-n+2k$ in $V_m \otimes V_n$ annihilated by Y. As the vectors from (6) form a basis of the $(-m-n+2k)$-weight subspace of $V_m \otimes V_n$, the latter problem reduces to the solution of the following system of linear equations with unknowns a_i's:

$$\begin{aligned}
0 &= Y\Big(\sum_{i=0}^{k} a_i\, e_{-m+2i} \otimes \tilde{e}_{-n+2(k-i)}\Big) \\
&= \sum_{i=0}^{k} a_i \left(Y e_{-m+2i} \otimes \tilde{e}_{-n+2(k-i)} + e_{-m+2i} \otimes Y \tilde{e}_{-n+2(k-i)}\right) \\
&= \sum_{i=1}^{k-1} \Big(a_i\, e_{-m+2(i-1)} \otimes \tilde{e}_{-n+2(k-i)} + a_i\, e_{-m+2i} \otimes \tilde{e}_{-n+2(k-i-1)}\Big) \\
&\qquad + a_0\, e_{-m} \otimes \tilde{e}_{-n+2(k-1)} + a_k\, e_{-m+2(k-1)} \otimes \tilde{e}_{-n} \\
&= \sum_{i=0}^{k-1} (a_{i+1} + a_i)\, e_{-m+2i} \otimes \tilde{e}_{-n+2(k-i-1)}.
\end{aligned}$$

This implies that $a_i = -a_{i+1}$ for all i, and the claim follows. ☐

To proceed we will need the following key result.

Lemma 3. *For all integers* k, l, m, n *with* $0 \leq l \leq k \leq \min\{m, n\}$, *the following identity holds*

$$\sum_{i=\max(0,2k-l-n)}^{\min(k,m-l)} (-1)^i \frac{(m-i)!\,(n-k+i)!}{i!\,(k-i)!\,(m-l-i)!\,(n+l-2k+i)!} = (-1)^{k+l}.$$

Proof. This identity can be restated as the Karlssor-Minton indentity for the hypergeometric series ${}_3F_2$. For a proof of the Karlsson-Minton identity, see [M]. ☐

To simplify the notation, denote $\mathfrak{s}_k = m + n - 2k$.

Lemma 4.

$$X^{\mathfrak{s}_k} b_{-\mathfrak{s}_k} = \mathfrak{s}_k! \sum_{l=0}^{k} (-1)^{k+l} \frac{(m-l)!\,(n-k+l)!}{l!\,(k-l)!} e_{m-2l} \otimes \tilde{e}_{n-2(k-l)}.$$

Proof. Direct calculation yields

$$\begin{aligned}
X^{\mathfrak{s}_k} b_{-\mathfrak{s}_k} &= X^{\mathfrak{s}_k} \Big(\sum_{i=0}^{k} (-1)^i e_{-m+2i} \otimes \tilde{e}_{-n+2(k-i)} \Big) \\
&= \sum_{j=0}^{n+m-2k} \sum_{i=0}^{k} (-1)^i \binom{m+n-2k}{j} (X^j e_{-m+2i}) \otimes \Big(X^{n+m-2k-j} \tilde{e}_{-n+2(k-i)} \Big) \\
&= \sum_{j=0}^{m} \sum_{i=m-k-j}^{m-j} (-1)^i \frac{(m+n-2k)!}{j!\,(m+n-2k-j)!} \times \\
&\qquad \times \frac{(m-i)!\,(j+i)!}{(m-i-j)!\,i!} \cdot \frac{(n-k+i)!\,(n+m-k-i-j)!}{(k+i+j-m)!\,(k-i)!} \times \\
&\qquad\quad \times e_{-m+2(i+j)} \otimes \tilde{e}_{-n+2(n+m-k-i-j)} \\
&= (\text{ substitution: } l = m - i - j\ ) \\
&= \mathfrak{s}_k! \sum_{l=0}^{k} \frac{(m-l)!\,(n-k+l)!}{l!\,(k-l)!} \times \\
&\qquad \times \Bigg(\sum_{i=\max\{0,2k-n-l\}}^{\min\{k,m-l\}} (-1)^i \frac{(m-i)!\,(n-k+i)!}{i!\,(k-i)!\,(m-l-i)!\,(n+l-2k+i)!} \Bigg) \times \\
&\qquad\quad \times e_{m-2l} \otimes \tilde{e}_{n-2(k-l)} \\
&= \mathfrak{s}_k! \sum_{l=0}^{k} (-1)^{k+l} \frac{(m-l)!\,(n-k+l)!}{l!\,(k-l)!} e_{m-2l} \otimes \tilde{e}_{n-2(k-l)},
\end{aligned}$$

where for the last equality we use Lemma 3. ☐

We are now ready to answer the Question from the introduction.

Theorem 5. *The forms ω_k are positive/negative, alternating with k, $0 \leq k \leq \min\{m, n\}$.*

Proof. Since $P_{-\mathfrak{s}_k}$ is spanned by $b_{-\mathfrak{s}_k}$, it suffices to verify that the value

$\omega_k(b_{-\mathfrak{s}_k}, b_{-\mathfrak{s}_k})$ is positive/negative, alternating with k.

$$\begin{aligned}
\omega_k(b_{-\mathfrak{s}_k}, b_{-\mathfrak{s}_k}) &= (Q \otimes R)(b_{-\mathfrak{s}_k}, X^{\mathfrak{s}_k}\, b_{-\mathfrak{s}_k}) \\
&= \mathfrak{s}_k! \sum_{i=0}^{k} \sum_{l=0}^{k} (-1)^{i+k+l} \frac{(m-l)!\,(n-k+l)!}{l!\,(k-l)!} \times \\
&\qquad \times Q(e_{-m+2i}, e_{m-2l}) R(\tilde{e}_{-n+2(k-i)}, \tilde{e}_{n-2(k-l)}) \\
&= \mathfrak{s}_k!\,(-1)^k qr \sum_{l=0}^{k} \frac{(m-l)!\,(n-k+l)!}{l!\,(k-l)!},
\end{aligned}$$

and the claim follows. □

Acknowledgements. I would like to thank Volodymyr Mazorchuk for all his help, and Svante Janson, Tobias Ekholm, Herbert Wilf, Doron Zeilberger, Christian Krattenthaler and Ganna Kudryavtseva for providing proofs for the identity of Lemma 3. I would also like to thank Hjalmar Rosengren and Michael Schlosser for pointing out that this identity is in fact the Karlsson-Minton identity for ${}_3F_2$. Finally, I would like to thank Anatoliy Klimyk for pointing out the connection to the problem of determining Clebsch-Gordan coefficients.

References

[BBFK] G. Barthel, J-P. Brasselet, K-H. Fieseler, L. Kaup, *Hodge-Riemann Relations for Polytopes — A Geometric Approach*, math.AG/0602411.

[FH] W. Fulton, J. Harris, *Representation theory. A first course*, Graduate Texts in Mathematics, 129. Readings in Mathematics, Springer-Verlag, New York, 1991.

[K] P. W. Karlsson, *Hypergeometric Functions with Integral Parameter Differences*, Journal of Mathematical Physics, Volume 12, 1971.

[M] B. M. Minton, Generalized Hypergeometric Function of Unit Argument, Journal of Mathematical Physics, Volume 11, 1970.

[PWZ] M. Petkovšek, H. S. Wilf, D. Zeilberger, $A = B$, With a foreword by Donald E. Knuth. A K Peters, Ltd., Wellesley, MA, 1996

[VK] N. Ja. Vilenkin, A. U. Klimyk, *Representation of Lie groups and special functions. Vol. 1. Simplest Lie groups, special functions and integral transforms*, Kluwer Academic Publishers Group, Dordrecht, 1991.

Department of Mathematics, Uppsala University, Box 480, SE-75106
Uppsala, Sweden, e-mail, `johan.kahrstrom@math.uu.se`.